

Applications of q-Calculus

Ali Omar

1. Introduction and Definition

Let ∇ denote the set of all analytic function $\hat{h}(z)$ in the open unit disk $\ddot{O} = \{z \in \mathbb{C}: |z| < 1\}$ with normalized condition $\hat{h}(0) = 0$ and $\hat{h}'(0) = 1$, and series expansion of $\hat{h}(z) \in \nabla$ is given as

$$\hat{h}(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

The star-like and convex functions $\hat{h} \in \nabla$ of order α can be defined as

$$\begin{aligned} \mathcal{S}^*(\alpha) &= \left\{ \hat{h} \in \nabla: \operatorname{Re} \left(\frac{z\hat{h}'(z)}{\hat{h}(z)} \right) > \alpha, 0 \leq \alpha < 1, z \in \ddot{O} \right\}, \\ \mathcal{C}(\alpha) &= \left\{ \hat{h} \in \nabla: \operatorname{Re} \left(\frac{(z\hat{h}'(z))'}{\hat{h}'(z)} \right) > \alpha, 0 \leq \alpha < 1, z \in \ddot{O} \right\}. \end{aligned}$$

Note that, if we take $\alpha = 0$, then the above defined classes of star-like and convex functions of order α convert into the well-known classes of star-like (δ^*) and convex (\mathcal{C}) functions. In 1991, the uniformly star-like functions (US) are introduced by Goodman [1] and defined as

$$\text{US} = \left\{ \hat{h} \in \nabla: \operatorname{Re} \left(\frac{z\hat{h}'(z)}{\hat{h}(z)} \right) > \left| \frac{z\hat{h}'(z)}{\hat{h}(z)} - 1 \right|, z \in \ddot{O} \right\}.$$

$$\begin{aligned} k - \text{UST} &= \{ \hat{h} \in \\ k - \text{UCV} &= \{ \hat{h} \in \\ &\text{as follows:} \\ &k^2((u-1)^2 + v^2) \\ &w-1 \mid \end{aligned}$$

Also, they defined these classes subject to the conic domain $\Omega_k(k \geq 0)$ (see [5]) as follows:

$$\begin{aligned} \Omega_k &= u + iv: u^2 > k^2((u-1)^2 + v^2), \\ &= w: \operatorname{Re} w > k|w-1| \end{aligned}$$

This domain represents the right-half plane for $k = 0$, hyperbola for $0 < k < 1$, parabola for $k = 1$, and ellipse $k > 1$. Furthermore, in [6], Shams et al. gave the generalization of k -uniformly star-like and k -uniformly convex functions and introduced a new subclasses $\text{DD}(k, \gamma)$ and $\text{SD}(k, \gamma)$ of analytic functions subject to the conic domain $\Omega_{k,\gamma}(k \geq 0)$, $0 \leq \gamma < 1$, as follows: After that, uniformly star-like functions (US) are studied by Ronning in [2] and Ma and Minda in [3].

Furthermore, in [4], Kanas and Wisniowska introduced the class k -uniformly star-like ($k - \text{UST}$) and k -uniformly convex functions ($k - \text{UCV}$), $k \geq 0$, as

$$\Omega_{k,\gamma} = \{u + iv: (u - \gamma)^2 > k^2((u - 1)^2 + v^2)\}.$$

$\Omega_{k,\gamma}$.

The function $p_{k,\gamma}(z)$ is extremal function for domain

$$p_{k,\gamma}(z) = \begin{cases} \Psi_1(z) & \text{for } k = 0 \\ \Psi_2(z) & \text{for } k = 1 \\ \Psi_3(z) & \text{for } 0 < k < 1 \\ \Psi_4(z) & \text{for } k > 1 \end{cases}$$

where

$$\begin{aligned} \Psi_1(z) &= \frac{1+z}{1-z}, \Psi_2(z) = 1 + \frac{2\gamma}{\pi^2} \left(\log \frac{1+\sqrt{z}}{1-\sqrt{z}} \right)^2 \\ \Psi_3(z) &= 1 + \frac{2\gamma}{1-k^2} \sinh^2 \left\{ \left(\frac{2}{\pi} \arccos k \right) \operatorname{arctanh} \sqrt{z} \right\} \\ \Psi_4(z) &= 1 + \frac{\gamma}{k^2-1} \sin \left(\frac{\pi}{2K(i)} \int_0^{u(z)/\sqrt{t}} \frac{1}{\sqrt{1-x^2}\sqrt{1-(ix)^2}} dx \right) + \frac{\gamma}{1-k^2}, \\ p_{k,\gamma}(z) &= 1 + Q_1 z + Q_2 z^2 + \dots, \end{aligned}$$

and $i \in (0,1)$; $k = \cosh \pi K'(i)/4K(i)$, $K(i)$ is the first kind of Legendre's complete elliptic integral. For details, see [5]. where Indeed, from (9), we have

$$\begin{aligned} Q_1 &= \begin{cases} \frac{2\gamma(2/\pi \arccos k)^2}{1-k^2} & \text{for } 0 \leq k < 1 \\ \frac{8\gamma}{\pi^2} & \text{for } k = 1 \\ \frac{\pi^2 \gamma}{4(1+t)\sqrt{t}K^2(t)(k^2-1)} & \text{for } k > 1 \end{cases} \\ Q_2 &= \begin{cases} \frac{(2/\pi \arccos k)^2 + 2}{3} Q_1 & \text{for } 0 \leq k < 1 \\ \frac{2}{3} Q_1 & \text{for } k = 1 \\ \frac{4K^2(t)(t^2 + 6t + 1) - \pi^2}{24K^2(t)(1+t)\sqrt{t}} Q_1 & \text{for } k > 1 \end{cases} \end{aligned}$$

as

The convolution of two functions \hat{h} and g can be defined

$$(\hat{h} * g)z = z + \sum_{n=2}^{\infty} a_n b_n z^n$$

where $\hat{h}(z)$ is given by

(1) and $g(z) = z + \sum_{n=2}^{\infty} b_n z^n, (z \in \tilde{O})$

If \hat{h} and $g \in \nabla$, we call that \hat{h} is subordinate to g , written as $\hat{h}(z) < g(z)$, such that $\hat{h}(z) = g(u(z))$, where $u(z) \in \nabla$ is the Schwarz function in \tilde{O} with conditions $u(0) = 0$ and $|u(z)| < 1$ (see [7])

Let P represent Carathéodory class of $p \in \nabla$, in the open unit disk \tilde{O} , and given by

$$p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$$

such that

$$\operatorname{Re}(p(z)) > 0.$$

In geometric function theory (GFT), the quantum (q -) calculus used as an important tools to study different families of analytic function and due to its application in mathematics and some related areas it has inspired the researcher very much. The quantum (or q^-) calculus is widely applied in various operators which include the q -difference (q -derivative) operator, and this operator plays an important role in GFT, quantum theory, number theory and statistical mechanics, etc. Jackson [8,9] was among the few researchers who defined the q -derivative and q -integral operator as well as provided some of their applications. Also, Ismail et al. [10] introduced research work in connection with function theory and q -theory. Later on, by using q -beta function, Gupta [11] introduced q -Baskakov-Durrmeyer operator, while q -Picard and q -Gauss-Weierstrass singular integral operators were introduced and studied by Aral, in [12]. Recently, Srivastava is one of the few researchers who studied univalent function theory by using a q -calculus (see, for details, [13,14]. Kanas and Raducanu [15] introduced Ruscheweyh q -differential operator, and Arif et al. [16] discussed some of its applications for multivalent functions. For more studies on q -analogous of operator, we refer [15,17 – 19]

$$[t]_q = 1 + q + q^2 + \cdots + q^{n-1} \quad (t = n \in \mathbb{N})$$

$$[n]_q! = \prod_{k=1}^n [k]_q, \quad (n \in \mathbb{N})$$

where $[0]_q! = 1$ and $[t]_q = 1 - q^t / 1 - q, t \in \mathbb{C}$.

$$[t]_{n,q} = \prod_{k=t}^{n-1} [k]_q, \quad (n \in \mathbb{N})$$

$$[t]_q = \frac{\Gamma_q(t+1)}{\Gamma_q(t)} \quad \text{and} \quad \Gamma_q(1) = 1$$

Definition 1 (see [6]). The q -number $[t]_q$ and q -factorial c for $q \in (0,1)$ is defined as

Definition 2. The q -generalized Pochhammer symbol and q -Gamma function be defined as

Definition 3 (see [9]). For $\hat{h} \in \nabla$, the q -derivative or (q -difference) operator be defined as

$$\partial_q \hat{h}(z) = \frac{-\hat{h}(z) + \hat{h}(qz)}{(q-1)z}, z \in \mathbb{D}$$

Combining (1) and (21), we have

$$\partial_q \hat{h}(z) = 1 + \sum_{n=2}^{\infty} [n]_q a_n z^{n-1}$$

Note that

$$\partial_q z^n = [n]_q z^{n-1}, \partial_q \left\{ \sum_{n=1}^{\infty} a_n z^n \right\} = \sum_{n=1}^{\infty} [n]_q a_n z^{n-1}$$

We can observe that

$$\lim_{q \rightarrow 1^-} \partial_q \hat{h}(z) = \hat{h}'(z).$$

Definition 4 (see [10]). Let an analytic function $\hat{h} \in \delta_q^*$ if

$$\left| \frac{z \partial_q \hat{h}(z)}{\hat{h}(z)} - \frac{1}{1-q} \right| \leq \frac{1}{1-q}$$

$$\hat{h}(0) = \hat{h}'(0) = 1$$

Equivalently, we can rewrite (26) and (25) as follows (see [20]) :

$$\frac{z \partial_q \hat{h}(z)}{\hat{h}(z)} < \frac{1+z}{1-qz}.$$

In [21], Zhang et al. defined the class $k - P_{q,\gamma}$ and generalized conic domain $\Omega_{k,q,\gamma}$.

Definition 5 (see [21]). Let $k \in [0, \infty)$, $q \in (0,1)$, and $\gamma \in \mathbb{C}/\{0\}$. A function $p(z) \in k - P_{q,\gamma}$ if and only if

$$p(z) < p_{k,\gamma,q}(z) = \frac{2p_{k,\gamma}(z)}{(1+q) + (1-q)p_{k,\gamma}(z)},$$

where $p_{k,\gamma}(z)$ is given by (9).

Geometrically, the function $p(z) \in k - P_{q,\gamma}$ takes all values from the domain $\Omega_{k,q,\gamma}$ which is defined as follows:

$$\Omega_{k,q,\gamma} = \gamma \Omega_{k,q} + (1-\gamma)$$

where

$$\Omega_{k,q} = \left\{ w: \operatorname{Re} \left(\frac{(1+q)w}{(q-1)w+2} \right) > k \left| \frac{(1+q)w}{(q-1)w+2} - 1 \right| \right\}$$

Remark 1. When $q \rightarrow 1 -$, then $\Omega_{k,q,\gamma} = \Omega_{k,\gamma}$, where $\Omega_{k,\gamma}$ is considered by Shams et al. [6].

Remark 2. When $\gamma = 1$ and $q \rightarrow 1 -$, then $\Omega_{k,q,\gamma} = \Omega_k$, where Ω_k is the conic domain considered by Kanas and Wisniowska [4].

Remark 3. For $\gamma = 1$ and $q \rightarrow 1 -$, then $k - P_{q,\gamma} = P(p_k)$, where $P(p_k)$ is the well-known class introduced by Kanas and Wisniowska [4].

Remark 4. For $\gamma = 1, k = 0$, and $q \rightarrow 1 -$, then $k - P_{q,\gamma} = P$, where P is the well-known Carathéodory class of analytic functions p

Mittag-Leffler introduced Mittag-Leffler function $\mathcal{H}_\alpha(z)$ in [22,23] as

$$\mathcal{H}_\alpha(z) = \sum_{n=0}^{\infty} \frac{1}{\Gamma(\alpha n + 1)} z^n, (\alpha \in \mathbb{C}, \operatorname{Re}(\alpha)) > 0,$$

and its generalization $\mathcal{H}_{\alpha,\beta}(z)$ is introduced by Wiman [24] as

$$\mathcal{H}_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{1}{\Gamma(\alpha n + \beta)} z^n, (\alpha, \beta \in \mathbb{C}, \operatorname{Re}(\alpha), \operatorname{Re}(\beta)) > 0.$$

For more study about Mittag-Leffler functions (see [25 – 30])

The q -Mittag-Leffler function defined by (see [31])

$$\mathcal{H}_{\alpha,\beta}(z, q) = \sum_{n=0}^{\infty} \frac{1}{\Gamma_q(\alpha n + \beta)} z^n, (\alpha, \beta \in \mathbb{C}, \operatorname{Re}(\alpha), \operatorname{Re}(\beta)) > 0$$

Note that q -Mittag-Leffler function is the specialized case of the q -Fox-Wright function $r\Phi_s(z, q)$ (see, for details, [32,33]). Since the q -Mittag-Leffler function $\mathcal{H}_{\alpha,\beta}(z, q)$ defined by (33) does not belong to the normalized analytic function class ∇ .

Now, we define the normalization of this q -Mittag-Leffler function $\mathcal{R}_{\alpha,\beta}(z)$ as

$$\begin{aligned} \mathcal{R}_{\alpha,\beta}(z, q) &= z \Gamma_q(\beta) \mathcal{H}_{\alpha,\beta}(z), \\ \mathcal{R}_{\alpha,\beta}(z, q) &= z + \sum_{n=2}^{\infty} \varphi_n(q, \beta) z^n, \\ \varphi_n(q, \beta) &= \frac{\Gamma_q(\beta)}{\Gamma_q(\alpha(n-1) + \beta)}, \end{aligned}$$

where $z \in \ddot{O}$, $\operatorname{Re} \alpha > 0$, and $\beta \in \mathbb{C}/\{0, -1, -2, \dots\}$. Corresponding to $\mathcal{R}_{\alpha,\beta}(z, q)$ and for $\hat{h} \in \nabla$, we define the following q -differential operator $\mathcal{D}_q^m(\alpha, \beta): \nabla \rightarrow \nabla$ by

$$\begin{aligned}
(\mathcal{D}_q^0(\alpha, \beta)\hat{h})z &= \hat{h}(z) * \mathcal{R}_{\alpha, \beta}(z, q) \\
(\mathcal{D}_q^1(\alpha, \beta)\hat{h})z &= z\mathcal{D}_q(\hat{h}(z) * \mathcal{R}_{\alpha, \beta}(z, q)) \\
(\mathcal{D}_q^2(\alpha, \beta)\hat{h})z &= \mathcal{D}_q(\mathcal{D}_q^1(\alpha, \beta)\hat{h})z \\
(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z &= \mathcal{D}_q(\mathcal{D}_q^{m-1}(\alpha, \beta)\hat{h})z
\end{aligned}$$

We note that

$$(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z = z + \sum_{n=2}^{\infty} [n]_q^m \varphi_n(q, \beta) a_n z^n$$

Note that,

- (i) For $\alpha = 0$ and $\beta = 1$, we have Salagean q -differential operator [34]
- (ii) For $q \rightarrow 1-$, $\alpha = 0$, and $\beta = 1$, we have Salagean differential operator [35]
- (iii) For $m = 0$, we have $E_{\alpha, \beta}(z, q)$ (see [31])
- (iv) For $m = 0$, we have $E_{\alpha, \beta}(z)$ (see [28])

We introduce a new subclass of analytic functions associated with newly defined q -differential operator (37).

Definition 6. Let $\hat{h}(z) \in \nabla$; then, $\hat{h}(z)$ is in the class $\mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$, $\gamma \in \mathbb{C}/\{0\}$, if it satisfies the condition

$$\operatorname{Re} \left\{ 1 + \frac{1}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{(\mathcal{D}_q^m((\alpha, \beta)\hat{h}))z} - 1 \right) \right\} > k \left| \frac{1}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{(\mathcal{D}_q^m((\alpha, \beta)\hat{h}))z} - 1 \right) \right|, \quad z \in \mathcal{O}$$

or equivalently

$$\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z} \in k - \mathcal{P}_{q, \gamma}.$$

Remark 5. For $\alpha = 0$ and $\beta = 1$, the class $\mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k) = k - \mathcal{US}(q, \gamma, m)$ studied by Hussain et al. [17].

Remark 6. For $m = 0$, $\alpha = 0$, $\beta = 1$, $q \rightarrow 1-$, and $\gamma = 1/1 - \eta$, $\eta \in \mathbb{C}/\{1\}$, the class $\mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k) = \mathcal{SD}(k, \eta)$ is studied by Shams et al. [6].

Remark 7. For $m = 0$, $\alpha = 0$, $\beta = 1$, $q \rightarrow 1-$ and $\gamma = 2/1 - \eta$, $\eta \in \mathbb{C}/\{1\}$, the class $\mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k) = \mathcal{DD}(k, \eta)$ is studied by Owa et al. [36].

Remark 8. For $k = 0$, $m = 0$, $\alpha = 0$, $\beta = 1$, $q \rightarrow 1-$, and $\gamma = 1/1 - \eta$, $\mathbb{C} \setminus \{1\}$, the class $\mathcal{US}_{\alpha, \beta}^{q, m}(\gamma, k) = \mathcal{S}^*(\eta)$ is the well-known subclass of star-like function.

Geometrically, an analytic function $\hat{h} \in \mathcal{U}\delta_{\alpha,\beta}^{q,m}(\gamma, k)$ if and only if

$$\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}$$

take all values in the generalized conic domain $\Omega_{k,q,\gamma}$ given by (29). Taking geometrical interpretation into consideration, one can rephrase the above definition as follows.

Definition 7. Let an analytic function $\hat{h} \in \mathcal{U}\delta_{\alpha,\beta}^{q,m}(\gamma, k)$ if and only if

$$\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} < p_{k,\gamma,q}(z),$$

where $p_{k,\gamma,q}(z)$ is given by (28).

2. Set of Lemmas

Lemma 1 (see [37]). If $F(z)$ is convex univalent in \tilde{O} and let $p(z) = \sum_{n=1}^{\infty} p_n z^n < F(z) = \sum_{n=1}^{\infty} d_n z^n$ in \tilde{O} , then

$$|p_n| \leq |d_1|, n \geq 1.$$

Lemma 2 (see [21]). Let $k \in [0, \infty)$ be fixed and

$$p_{k,\gamma,q}(z) = \frac{2p_{k,\gamma}(z)}{(1+q) + (1-q)p_{k,\gamma}(z)}.$$

Then,

$$p_{k,\gamma,q}(z) = 1 + \frac{2}{1+q} Q_1 z + \left\{ \frac{2}{1+q} Q_2 - \frac{2(1-q)}{1+q} Q_1^2 \right\} z^2 + \dots,$$

where Q_1 and Q_2 are given by (12) and (13).

Lemma 3 (see [21]). Let $p(z) = 1 + \sum_{n=1}^{\infty} p_n z^n \in k - P_{q,\gamma}$; then,

$$|p_n| \leq \frac{2}{1+q} |Q_1|, n \geq 1.$$

Lemma 4 (see [38]). Let an analytic function $p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n \in P$ and $\text{Re}(p(z)) > 0$ for all z in \tilde{O} ; then,

$$|c_2 - \mu c_1^2| \leq 2 \max\{1, |2\mu - 1|\}, \forall \mu \in \mathbb{C}.$$

3. Main Results

Theorem 1. Let an analytic function $\hat{h}(z) \in \mathcal{U}\delta_{\alpha,\beta}^{q,m}(\gamma, k)$; then,

$$(\mathcal{D}_q^m((\alpha, \beta)\hat{h})z) < z \exp \int_0^z \frac{p_{k,\gamma,q}(u(\xi)) - 1}{\xi} d\xi,$$

where $u(z)$ is analytic in \ddot{O} having condition $u(0) = 0$ and $|u(z)| < 1$. Furthermore, for $|z| = \rho$, we have

$$\begin{aligned} \exp \int_0^1 \frac{p_{k,\gamma,q}(-\rho) - 1}{\rho} d\rho &\leq \left| \frac{(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{z} \right| \\ &\leq \exp \int_0^1 \frac{p_{k,\gamma,q}(\rho) - 1}{\rho} d\rho, \end{aligned}$$

where $p_{k,\gamma,q}(z)$ is defined by (28).

Proof. If $\hat{h}(z) \in \mathcal{U}\delta_{\alpha,\beta}^{q,m}(\gamma, k)$, then, from identity (41), we have

$$\begin{aligned} \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z} &= p_{k,\gamma,q}(u(z)), \\ \frac{\partial_q (\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z} - \frac{1}{z} &= \frac{p_{k,\gamma,q}(u(z)) - 1}{z}. \end{aligned}$$

Integrating (50) and after some simplification, we have

$$(\mathcal{D}_q^m((\alpha, \beta)\hat{h})z) < z \exp \int_0^z \frac{p_{k,\gamma,q}(u(\xi)) - 1}{\xi} d\xi.$$

Hence, (47) is proved. Noting that the univalent function $p_{k,\gamma,q}(z)$ maps the disk $|z| < \rho$ ($0 < \rho \leq 1$) onto a region which is convex and symmetric with respect to the real axis, we see

$$\begin{aligned} p_{k,\gamma,q}(-\rho|z|) &\leq \operatorname{Re}\{p_{k,\gamma,q}(u(\rho z))\} \\ &\leq p_{k,\gamma,q}(\rho|z|) \quad (0 < \rho \leq 1, z \in \ddot{O}). \end{aligned}$$

Using (51) and (52) gives

$$\int_0^1 \frac{p_{k,\gamma,q}(-\rho|z|) - 1}{\rho} d\rho \leq \operatorname{Re} \int_0^1 \frac{p_{k,\gamma,q}(u(\rho(z))) - 1}{\rho} d\rho \leq \int_0^1 \frac{p_{k,\gamma,q}(\rho|z|) - 1}{\rho} d\rho,$$

for $z \in \ddot{O}$. Thus, inequality (51) leads us to

$$\begin{aligned} \int_0^1 \frac{p_{k,\gamma,q}(-\rho|z|) - 1}{\rho} d\rho &\leq \log \left| \frac{(\mathcal{D}_q^m(\alpha, \beta)\hat{h})z}{z} \right| \leq \int_0^1 \frac{p_{k,\gamma,q}(\rho|z|) - 1}{\rho} d\rho, \\ p_{k,\gamma,q}(-\rho) &\leq p_{k,\gamma,q}(-\rho|z|), p_{k,\gamma,q}(\rho|z|) \leq p_{k,\gamma,q}(\rho), \end{aligned}$$

which implies that

$$\begin{aligned} \exp \int_0^1 \frac{p_{k,\gamma,q}(-\rho) - 1}{\rho} d\rho &\leq \left| \frac{(\mathcal{D}_q^m(\alpha, \beta) \hat{h})z}{z} \right| \\ &\leq \exp \int_0^1 \frac{p_{k,\gamma,q}(\rho) - 1}{\rho} d\rho \end{aligned}$$

Hence, the proof is complete.

For $\alpha = 0$ and $\beta = 1$, then we have the following known corollary proved by Hussain et al. in [17].

Corollary 1. Let an analytic function $\hat{h}(z) \in k\text{-US}(q, \gamma, m)$; then,

$$\mathcal{D}_q^m \hat{h}(z) < z \exp \int_0^z \frac{p_{k,\gamma}(u(\xi)) - 1}{\xi} d\xi,$$

where $u(z)$ is analytic in \tilde{O} having condition $u(0) = 0$ and $|u(z)| < 1$. Furthermore, for $|z| = \rho$, we have

$$\exp \int_0^1 \frac{p_{k,\gamma}(-\rho) - 1}{\rho} d\rho \leq \left| \frac{\mathcal{D}_q^m \hat{h}(z)}{z} \right| \leq \exp \int_0^1 \frac{p_{k,\gamma}(\rho) - 1}{\rho} d\rho. \text{ where } p_{k,\gamma}(z) \text{ is defined by (9).}$$

Theorem 2. If $\hat{h}(z) \in \mathcal{U}\delta_{\alpha,\beta}^{q,m}(\gamma, k)$, then

$$\begin{aligned} |a_2| &\leq \frac{\delta}{[2]_q^m \{ [2]_q - 1 \} \varphi_2(q, \beta)}, \\ |a_n| &\leq \frac{\delta}{[n]_q^m \{ [n]_q - 1 \} \varphi_n(q, \beta)} \\ &\quad \cdot \prod_{j=1}^{n-2} \left(1 + \frac{\delta}{[j+1]_q - 1} \right), \text{ for } n \geq 3, \end{aligned}$$

where $\delta = 2|Q_1|/1 + q$ with Q_1 and $\varphi_n(q, \beta)$ is given by (12) and (35).

Proof. Let

$$\frac{z \partial_d (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} = p(z),$$

where $p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$ and $(\mathcal{D}_q^m(\alpha, \beta) \hat{h})z$ is given by (37); then, (61) becomes

$$z + \sum_{n=2}^{\infty} [n]_q^{m+1} \varphi_n(q, \beta) a_n z^n = \left(\sum_{n=0}^{\infty} c_n z^n \right) \left(z + \sum_{n=2}^{\infty} [n]_q^m \varphi_n(q, \beta) a_n z^n \right)$$

Now, comparing the coefficients of z^n , we obtain

$$[n]_q^{m+1} \varphi_n(q, \beta) a_n = [n]_q^m \varphi_n(q, \beta) a_n + \sum_{j=1}^{n-1} [j]_q^m \varphi_j(q, \beta) a_j c_{n-j}$$

which implies

$$a_n = \frac{1}{[n]_q^m \{[n]_q - 1\} \varphi_n(q, \beta)} \sum_{j=1}^{n-1} [j]_q^m \varphi_j(q, \beta) a_j c_{n-j}$$

Using Lemma 4, we have $|a_n| \leq \frac{2|Q_1|}{(1+q)[n]_q^m \{[n]_q - 1\} \varphi_n(q, \beta)} \sum_{j=1}^{n-1} [j]_q^m \varphi_j(q, \beta) |a_j|$

Let us take $\delta = 2|Q_1|/1 + q$. Then, we have

$$|a_n| \leq \frac{\delta}{[n]_q^m \{[n]_q - 1\} \varphi_n(q, \beta)} \sum_{j=1}^{n-1} [j]_q^m \varphi_j(q, \beta) |a_j|$$

For $n = 2$, in (66), we have

$$|a_2| \leq \frac{\delta}{[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)}$$

which shows that (60) holds for $n = 2$. To prove (60), we use the principle of mathematical induction; for this, consider the case $n = 3$:

$$|a_3| \leq \frac{\delta}{[3]_q^m \{[3]_q - 1\} \varphi_3(q, \beta)} \{1 + [2]_q^m \varphi_2(q, \beta) |a_2|\}$$

Using (67), we have

$$|a_3| \leq \frac{\delta}{[3]_q^m \{[3]_q - 1\} \varphi_3(q, \beta)} \left\{ 1 + \frac{\delta}{[2]_q - 1} \right\}$$

which shows that (60) holds for $n = 3$. Let us assume that (60) is true for $n \leq t$, that is,

$$|a_t| \leq \frac{\delta}{[t]_q^m \{[t]_q - 1\} \varphi_t(q, \beta)} \prod_{j=1}^{t-2} \left(1 + \frac{\delta}{[j+1]_q - 1} \right), \text{ for } n = 3, 4, \dots$$

Consider

$$\begin{aligned}
|a_{t+1}| &\leq \frac{\delta}{[t+1]_q^m \{[t+1]_q - 1\} \varphi_{t+1}(q, \beta)} \\
&\leq \frac{\delta}{[t+1]_q^m \{[t+1]_q - 1\} \varphi_{t+1}(q, \beta)} \\
&\quad \times \left\{ 1 + [2]_q^m \varphi_2(q, \beta) |a_2| + [3]_q^m \varphi_3(q, \beta) |a_3| \right. \\
&\quad \left. + [4]_q^m \varphi_4(q, \beta) |a_4| + \cdots + [t]_q^m \varphi_t(q, \beta) |a_t| \right\} \\
&\leq \frac{\delta}{[t+1]_q^m \{[t+1]_q - 1\} \varphi_{t+1}(q, \beta)} \\
&\quad \times \left\{ 1 + \frac{\delta}{[2]_q - 1} + \frac{\delta}{[3]_q - 1} \left(1 + \frac{\delta}{[2]_q - 1} \right) \right. \\
&\quad \left. + \cdots + \frac{\delta}{[t]_q - 1} \prod_{j=1}^{t-2} \left(1 + \frac{\delta}{[j+1]_q - 1} \right) \right\} \\
&= \frac{\delta}{[t+1]_q^m \{[t+1]_q - 1\} \varphi_{t+1}(q, \beta)} \prod_{j=1}^{t-1} \left(1 + \frac{\delta}{[j+1]_q - 1} \right)
\end{aligned}$$

which shows that (60) holds for $n = t + 1$. Hence, theorem is completed.

For $\alpha = 0$ and $\beta = 1$, then we have the following known corollary proved by Hussain et al., in [17].

Corollary 2. If $\hat{h}(z) \in \mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$, then

$$\begin{aligned}
|a_2| &\leq \frac{\delta}{[2]_q^m \{[2]_q - 1\} \varphi_2} \\
|a_n| &\leq \frac{\delta}{[n]_q^m \{[n]_q - 1\}} \prod_{j=1}^{n-2} \left(1 + \frac{\delta}{[j+1]_q - 1} \right), \text{ for } n \geq 3
\end{aligned}$$

where $\delta = 2|Q_1|/1 + q$ and Q_1 is given by (12).

Theorem 3. Let $\hat{h}(z) \in \mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$ of form (1) and $0 \leq k < \infty$ be fixed and $\mu \in \mathbb{C}$; then,

$$|a_3 - \mu a_2^2| \leq \frac{Q_1}{[3]_q^m \varphi_3(q, \beta) (1 + q) \{[3]_q - 1\}}^m [1, |2v - 1|],$$

where

$$v = \frac{1}{2} \left(1 - \frac{Q_2}{Q_1} + (1 - q)Q_1 - \frac{2Q_1}{(1 + q)\{[2]_q - 1\}} \right. \\
\left. + \mu \frac{Q_1 [3]_q^m \{[3]_q - 1\}}{\varphi_2(q, \beta) (1 + q) ([2]_q^m \{[2]_q - 1\})^2} \right)$$

where Q_1 and Q_2 are given by (12) and (13).

Proof. Let $\hat{h}(z) \in \mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$; then, from (12), we have

$$\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} = p_{k, \gamma}(u(z)) z \in \ddot{O}$$

Let

$$p(z) = \frac{1}{(1 - u(z))(1 + u(z))^{-1}} = 1 + c_1 z + c_2 z^2 + \dots$$

and this gives

$$\begin{aligned} u(z) &= \frac{c_1}{2} z + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots \\ &= 1 + \frac{Q_1 c_1}{(1+q)} z + \frac{1}{(1+q)} \left\{ \frac{Q_2 c_1^2}{2} + \left(c_2 - \frac{c_1^2}{2} \right) Q_1 \right. \\ &\quad \left. - \frac{(1-q) Q_1^2 c_1^2}{2} \right\} z^2 + \dots, \end{aligned}$$

$$\begin{aligned} &\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} \\ &= 1 + [2]_q^m \varphi_2(q, \beta) \{ [2]_q - 1 \} a_2 z \\ &\quad + \{ [3]_q^m \varphi_3(q, \beta) \{ [3]_q - 1 \} a_3 \\ &\quad - ([2]_q^m \varphi_2(q, \beta))^2 \{ [2]_q - 1 \} a_2^2 \} z^2. \end{aligned}$$

Using (79) in (76) and comparing with (80), we obtain

Theorem 4. If an analytic function $\hat{h}(z)$ ofform (1) satisfies the condition,

$$\sum_{n=2}^{\infty} \left\{ \{ [n]_q - 1 \} (k+1) + |\gamma| \right\} |\varphi_n(q, \beta)| | [n]_q^m | |a_n| \leq |\gamma|$$

then $\hat{h}(z) \in \mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$

Proof. Let we note that

$$\begin{aligned} \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right| &= \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z - (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} \right| \\ &= \frac{\sum_{n=2}^{\infty} [n]_q^m \varphi_n(q, \beta) \{ [n]_q - 1 \} a_n z^n}{z + \sum_{n=2}^{\infty} [n]_q^m \varphi_n(q, \beta) a_n z^n} \\ &\leq \frac{\sum_{n=2}^{\infty} | [n]_q^m \varphi_n(q, \beta) \{ [n]_q - 1 \} | |a_n|}{1 - \sum_{n=2}^{\infty} | [n]_q^m | |\varphi_n(q, \beta)| \| a_n \|}. \end{aligned}$$

From (85), it follows that

$$1 - \sum_{n=2}^{\infty} |[n]_q^m| |\varphi_n(q, \beta)| |a_n| > 0.$$

To show that $\hat{h}(z) \in \mathcal{US}_{\alpha, \beta}^{q, m}(\gamma, k)$, it suffices that

$$a_2 = \frac{Q_1 c_1}{[2]_q^m \varphi_2(q, \beta)(1+q)\{[2]_q - 1\}}$$

$$a_3 = \frac{1}{[3]_q^m \varphi_3(q, \beta)\{[3]_q - 1\}} \times \left\{ \frac{Q_1 c_2 / 1 + q}{+ \frac{1}{1+q} \left(\frac{Q_2}{2} - \frac{Q_1}{2} - \frac{(1-q)Q_1^2}{2} + \frac{Q_1^2}{(1+q)\{[2]_q - 1\}} \right) c_1^2} \right\},$$

$$a_3 - \mu a_2^2 = \frac{Q_1}{[3]_q^m \varphi_3(q, \beta)\{[3]_q - 1\}(1+q)}$$

$$\times \left\{ c_2 - \frac{1}{2} \left(1 - \frac{Q_2}{Q_1} + (1-q)Q_1 - \frac{2Q_1}{(1+q)\{[2]_q - 1\}} \right) + \mu \frac{Q_1 [3]_q^m \{[3]_q - 1\}}{\varphi_3(q, \beta)(1+q)([2]_q^m \{[2]_q - 1\})^2} \right\} c_1^2.$$

For any complex number μ and after some calculation, we have

$$a_3 - \mu a_2^2 = \frac{Q_1}{[3]_q^m \varphi_3(q, \beta)(1+q)\{[3]_q - 1\}} \{c_2 - v c_1^2\}$$

where v is given by (75). Using Lemma 4 on (82), we have the required results.

For $\alpha = 0$ and $\beta = 1$, then we have the following known corollary proved by Hussain et al. in [17].

Corollary 3. Let $\hat{h}(z) \in \mathcal{US}_{\alpha, \beta}^{q, m}(\gamma, k)$ of form (1) and $0 \leq k < \infty$ be fixed and $\mu \in \mathbb{C}$; then,

$$|a_3 - \mu a_2^2| \leq \frac{Q_1}{2[3]_q^m \{[3]_q - 1\}} m [1, |2v - 1|]$$

where

$$= \frac{1}{2} \left(1 - \frac{Q_2}{Q_1} + (1-q)Q_1 - \frac{Q_1}{\{[2]_q - 1\}} + \mu \frac{Q_1 [3]_q^m \{[3]_q - 1\}}{2([2]_q^m \{[2]_q - 1\})^2} \right)$$

where Q_1 and Q_2 are given by (12) and (13), where $p_{k,\gamma}(z)$ is defined by (9)).

$$\left| \frac{k}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right) \right| - \operatorname{Re} \left\{ \frac{1}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right) \right\} \leq 1$$

From (86), we have

$$\begin{aligned} & \leq \left| \frac{k}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right) \right| - \operatorname{Re} \left\{ \frac{1}{\gamma} \left(\frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right) \right\} \\ & \leq \frac{k}{|\gamma|} \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right| + \frac{1}{|\gamma|} \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right| \\ & \leq \frac{(k+1)}{|\gamma|} \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} - 1 \right| \\ & = \frac{(k+1)}{|\gamma|} \left| \frac{z \partial_q (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z - (\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z}{(\mathcal{D}_q^m(\alpha, \beta) \hat{h}) z} \right| \\ & \leq \frac{(k+1)}{|\gamma|} \left\{ \frac{|\sum_{n=2}^{\infty} [n]_q^m \varphi_n(q, \beta) \{[n]_q - 1\}| |a_n|}{1 - \sum_{n=2}^{\infty} |[n]_q^m| |\varphi_n(q, \beta)| |a_n|} \right\} 1, \end{aligned}$$

from (85)

When $q \rightarrow 1-, m = 0, \alpha = 0, \beta = 1$, and $\gamma = 1 - \eta$, with $0 \leq \eta < 1$, then we have the following.

Corollary 4 (see [6]). An analytic function \hat{h} of form (1) belongs to the class $k - \text{US}(1 - 2\eta)$ if it holds the following condition:

$$\sum_{n=2}^{\infty} \{n(k+1) - (k+\eta)\} |a_n| \leq 1 - \eta$$

where $0 \leq \eta < 1$ and $k \geq 0$.

When $q \rightarrow 1-, m = 0, \alpha = 0, \beta = 1$, and $\gamma = 1 - \eta$, with $0 \leq \eta < 1$ and $k = 0$, then we have the following.

Corollary 5 (see [39]). An analytic function $\hat{h} \in \text{US}(1 - \eta)$ of form (1) it holds

$$\sum_{n=2}^{\infty} \{n - \eta\} |a_n| \leq 1 - \delta, 0 \leq \eta < 1.$$

Theorem 5. Let $\hat{h}(z) \in \mathcal{U}_{\alpha, \beta}^{q, m}(\gamma, k)$; then, $\hat{h}(\ddot{O})$ contains an open disk of radius:

$$\frac{[2]^m \varphi_2(q, \beta) \{[2]_q - 1\}}{(2[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)) + \delta}$$

where $\delta = 2|Q_1|/1 + q$ with Q_1 is given by (12). Proof. Let $u_0 \neq 0$ be a complex number such that $\hat{h}(z) \neq u_0$ for $z \in \tilde{O}$. Then,

$$\hat{h}_1(z) = \frac{u_0 \hat{h}(z)}{u_0 - \hat{h}(z)} = z + \left(a_2 + \frac{1}{u_0}\right)z^2 + \dots$$

Since $\hat{h}_1(z)$ is univalent, so

$$\left|a_2 + \frac{1}{u_0}\right| \leq 2.$$

Now, using (59), we have

$$\left|\frac{1}{u_0}\right| \leq \frac{(2[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)) + \delta}{[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)}$$

Hence, we have

$$|u_0| \geq \frac{[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)}{(2[2]_q^m \{[2]_q - 1\} \varphi_2(q, \beta)) + \delta}$$

For $\alpha = 0$ and $\beta = 1$, then we have the following known corollary proved by Hussain et al. in [17].

Corollary 6. Let $\hat{h}(z) \in \mathcal{US}_{\alpha, \beta}^{q, m}(\gamma, k)$; then, $\hat{h}(\tilde{O})$ contains an open disk of radius:

$$\frac{[2]_q^m \{[2]_q - 1\}}{2[2]_q^m \{[2]_q - 1\} + \delta}$$

where $p_{k, \gamma}(z)$ is defined by (9) and $\delta = |Q_1|$ with Q_1 is given by (12).

4. Conclusion

Here, we have introduced and defined new subclass of analytic functions in generalized conic domain by using newly defined q -differential operator. This newly defined operator for analytic functions is the extension of q -Salagean operator. We investigated structural formula, coefficient estimates, and Fekete-Szegő problem, and also, we have successfully derived subordination result.

.

8. References

1. M. Abramowitz and I. A. Stegun (Eds.), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Dover Publications Inc., New York, 1965.

2. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, *Filomat* 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
3. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, *Complex Var. Elliptic Equ.* 61(3) (2016), 338-350.
<https://doi.org/10.1080/17476933.2015.1079628>
4. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, *Commun. Korean Math. Soc.* 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
5. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, *Fract. Calc. Appl. Anal.* 5(3) (2002), 303-313.
6. P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften Series 259, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, *Kyungpook Math. J.* **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>
7. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, *Konuralp J. Math.* 7(1) (2019), 199-202.
8. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, *Stud. Univ. Babeş-Bolyai Math.* 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbmath.2020.1.06>
9. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, *Afr. Mat.* 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
10. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, *J. Appl. Math.* 2011(2011), Article ID 298628.
<https://doi.org/10.1155/2011/298628>
11. V. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.
12. E. Merkes and B. T. Scott, Starlike hypergeometric functions, *Proc. Amer. Math. Soc.* 12 (1961), 885-888.

13. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, C. R. Acad. Sci. Paris 137(1903), 554 – 558.
14. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, Integral Transforms Spec. Funct. 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
15. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, Novi Sad J. Math. 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
16. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, Hacet. J. Math. Stat. 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS> . 20164513110
17. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, Afr. Mat. 28(2017), 1357-1366.
<https://doi.org/10.1007/s13370-017-0520-x>
18. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, Afr. Mat. 27(5) (2016), 1021-1027.
<https://doi.org/10.1007/s13370-016-0398-z>
19. M. S. Robertson, On the theory of univalent functions, Ann. of Math. (2) 37(2) (1936), 374-408.
20. H. Silverman, Starlike and convexity properties for hypergeometric functions, J. Math. Anal. Appl. 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
21. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, Integral Transforms Spec. Funct. 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
22. L. Spáček, Contribution à la théorie des fonctions univalentes, Časopis Pro Pěstování Matematiky 62(1932), 12 – 19.
23. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, Journal of Inequalities in Pure and Applied Mathematics 5(4) (2004), Article ID 83, 10 pages.
24. A. Wiman, Über die nullstellen der funktionen $E_\alpha(x)$, Acta Math. 29 (1905), 217-134.

25. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, Yokohama Math. J. 19 (1971), 7 – 15.
26. Salah, J. and Darus, M., A note on Generalized Mittag-Leffler function and Application, Far East Journal of Mathematical Sciences (FJMS) 48 (1), 33–46, 2011.
27. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein](#) [AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, Progr. Fract. Differ. Appl. Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL: <https://www.naturalspublishing.com/Article.asp?ArtcID=26373>
28. Jamal Salah, Maryam Al Hashmi and Khaled Al Mashrafi, Some propositions on mathematical models of population growth, International Journal of Latest Trends in Engineering and Technology (IJLTET)Special Issue - ICTIMESH - 2022, 64-69.
URL: https://www.ijltet.org/journal_details.php?id=978&j_id=5050
29. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Inclusion Results of a Generalized Mittag-Leffler-Type Poisson Distribution in the k-Uniformly Janowski Starlike and the k-Janowski Convex Functions, Mathematics and Statistics, Vol. 11, No. 1, pp. 22-27, 2023.
DOI: 10.13189/ms.2023.110103
30. Jamal Salah, [Riemann and Big Bang Hypotheses](#), NEUROQUANTOLOGY, Vol. 20, No. 16, pp. 1592-1597, 2022. DOI: 10.48047/NQ.2022.20.16. NQ880156
31. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Subclasses of Spiral-Like Functions Associated with the Generalized Mittag-Leffler Function, NEUROQUANTOLOGY, Vol. 20, No. 15, pp. 4784-4799, 2022. DOI: 10.14704/NQ.2022.20.15. NQ88484
32. Jamal Salah, Maryam Al Hashmi, Hameed Ur Rehman, Khaled Al Mashrafi, Modified Mathematical Models in Biology by the Means of Caputo Derivative of A Function with Respect to Another Exponential Function, Mathematics and Statistics, Vol. 10, No. 6, pp. 1194 - 1205, 2022. DOI: 10.13189/ms.2022.100605
33. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function, Mathematics and Statistics, vol. 10, no. 2, pp. 410-418, 2022, DOI: 10.13189/ms.2022.100216.
34. Jamal Salah, Some Remarks and Propositions on Riemann Hypothesis, Mathematics and Statistics, vol. 9, no. 2, pp. 159-165, 2021, DOI: 10.13189/ms.2021.090210.
35. Hameed Ur Rehman, Malina Darus, and Jamal Salah, Generalizing Certain Analytic Functions Correlative to the n-th Coefficient of Certain Class of Bi-Univalent Functions, Hindawi Journal of Mathematics Volume 2021, Article ID 6621315, 14 pages DOI: [10.1155/2021/6621315](https://doi.org/10.1155/2021/6621315)

36. Salah, J., (2021). Applications of differential Subordination and Superordination Involving Certain Fractional Operator. Current Topics on Mathematics and Computer Science, vol. 1, pp. 7–16, 2021 [DOI: 10.9734/bpi/ctmcs/v1/9050D](https://doi.org/10.9734/bpi/ctmcs/v1/9050D)
37. Jamal Y. Mohammad Salah, The Consequence of the Analytic Continuity of Zeta Function Subject to an Additional Term and a Justification of the Location of the Non-Trivial Zeros, International Journal of Science and Research (IJSR), Vol. 9, Issue 3, March 2020, pp. 1566-1569, March 2020, URL: https://www.ijsr.net/get_abstract.php?paper_id=SR20328201050
38. Jamal Y. Mohammad Salah, An Alternative perspective to Riemann Hypothesis, PSYCHOLOGY AND EDUCATION, vol. 57, no. 9, pp. 1278-1281, 2020, URL: [file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20\(1\).pdf](file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20(1).pdf)
39. Jamal Y. Mohammad Salah, Two Conditional proofs of Riemann Hypothesis, International Journal of Sciences: Basic and Applied Research (IJSBAR), vol. 49, no. 1, pp. 74-83, 2020, URL: <https://gssrr.org/index.php/JournalOfBasicAndApplied/article/view/10720>
40. Jamal Salah, TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS, Far East Journal of Mathematical Sciences (FJMS) Volume 118, Issue1, 2019, Pages 1- 8. Sept 2019.
DOI: 10.17654/MS118010001
41. Jamal Y. Salah, A NEW SUBCLASS OF UNIVALENT FUNCTIONS DEFINED BY THE MEANS OF JAMAL OPERATOR, Far East Journal of Mathematical Sciences (FJMS) Vol. 108, No. 2, 2018, pp. 389-399, 2018, Dec 2018, DOI: 10.17654/MS108020389.
42. Hameed Ur Rehman, Maslina Darus and **Jamal Salah**, Graphing Examples of Starlike and Convex Functions of order β , Appl. Math. Inf. Sci. vol. 12, No. 3, pp. 509-515, 2018, DOI:10.18576/amis/120305
43. Jamal Salah, Hameed Ur Rehman and Maslina Darus, A Note on Caputo's derivative interpretation in Economy, J. Appl. Math, pp. 1-7, 2018, DOI: 10.1155/2018/1260240.
44. Hameed Ur Rehman, Maslina Darus and Jamal Salah, Normalization of the generalized K– Mittag-Leffler function and ratio to its sequence of partial sums, BSG Proceedings, vol. 25, 2018, pp. 78-94.
45. Hameed Ur Rehman, Maslina Darus and Jamal Salah, COEFFICIENT PROPERTIES INVOLVING THE GENERALIZED K–MITTAG-LEFFLER FUNCTIONS, Trans. J. Math. Mecha. (TJMM) 9 (2017), No. 2, 155-164
46. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
47. Jamal Y. Salah. Properties of the Modified Caputo's Derivative Operator for certain analytic functions. International Journal of Pure and Applied Mathematics. September 2016, Vol. 109, No. 3, pp. 665 – 671, 2014, DOI: 10.12732/ijpam.v109i3.14
48. Jamal Salah. A Note on the Modified Caputo's Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 100, No. 4, pp. 609-615, 2016,
DOI: 10.17654/MS100040609
49. S. Venkatesh, Jamal Salah, G. Sethuraman. "Some Results on E – Cordial Graphs", *International Journal of Mathematical Trends and Technology (IJMTT)*. V7:121-125 March 2014.
DOI: 10.14445/22315373/IJMTT-V7P516

50. Jamal Y. Salah, A Note on Gamma Function, International Journal of Modern Sciences and Engineering Technology (IJMSET), vol. 2, no. 8, pp. 58-64, 2015,
51. Jamal Y. Salah, On Riemann Hypothesis and Robin's Inequality, International Journal of Scientific and Innovative Mathematical Research (IJSIMR), vol. 3, no. 4, pp. 9-14, April 2015,
URL: <https://www.arcjournals.org/pdfs/ijsimr/v3-i4/4.pdf>
52. Jamal Y. Salah, **A Note on Riemann Zeta Function**, IOSR Journal of Engineering (IOSRJEN), vol. 06, no. 02, pp. 07-16, February 2016,
URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
53. Jamal Y. Salah, A NOTE ON THE HURWITZ ZETA FUNCTION, Far East Journal of Mathematical Sciences (FJMS), vol. 101, no. 12, pp. 2677-2683, June 2017,
DOI: [10.17654/MS101122677](https://doi.org/10.17654/MS101122677)
54. Jamal Salah and Venkatesh Srivastava. Inequalities on the Theory of Univalent Function. Journal of Mathematics and System Science, vol. 4, pp. 509-513, 2014.

DOI: 10.17265/2159-5291/2014.07.008
55. Jamal Salah. Neighborhood of a certain family of multivalent functions with negative coefficients. International Journal of Pure and Applied Mathematics. (IJPAM). Vol. 92, No 4, April 2014,
DOI: 10.12732/ijpam.v92i4.14
56. Jamal Salah. Fekete-Szego problems involving certain integral operator. International Journal of Mathematics Trends and Technology. IJMTT, vol. 7, no. 1, pp. 54-60, 2014.
DOI: 10.14445/22315373/IJMTT-V7P508
57. Jamal Salah. Subordination and superordination involving certain fractional operator. Asian Journal of Fuzzy and Applied Mathematics, vol. 1, pp. 98-107, 2013.
URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>
58. Jamal Salah. Certain subclass of analytic functions associated with fractional calculus operator. Trans. J. Math. Mecha, vol. 3, no. 1, pp. 35-42, 2011.
URL: <http://tjmm.edyropress.ro/journal/11030106.pdf>
59. Jamal Salah. A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional. J. Appl. comp Sc. Math, vol. 5, no. 1, pp. 97- 101, 2011,
URL: https://www.jacsm.ro/view/?pid=10_16
60. Jamal Salah and Maslina Darus. On convexity of general integral operators.
J. Anale. Universitatii. De Vest. Timisoara. Seria
Matematica – Informatica XLIX. 1 (2011) 117-124.
61. Jamal Salah and Maslina Darus. A note on generalized Mittag-Leffler and Application. Far East. Math Sc. (FJMS), vol. 48, no. 1, pp. 33-46, 2011.
URL: <http://www.pphmj.com/abstract/5478.htm>
62. Jamal Salah and Maslina Darus. A subclass of uniformly convex functions associated with fractional calculus operator involving Caputo's fractional differentiation. Acta. Univ. APL. Vol. 24, pp. 295-306, 2010.

63. D. G. Cantor, "Power series with integral coefficients," *Bulletin of the American Mathematical Society*, vol. 69 , no. 3, pp. 362 – 367, 1963.
64. R. Vein and P. Dale, "Determinants and Their Applications in Mathematical Physics," in *Applied Mathematical Sciences*, vol. 134, Springer, New York, 1999.
65. R. Wilson, "Determinantal criteria for meromorphic functions," *Proceedings of the London Mathematical Society*, vol. s3-4, no. 1, pp. 357-374, 1954.
66. C. Pommerenke, "On the coefficients and Hankel determinants of univalent functions," *Journal of the London Mathematical Society*, vol. s1-41, no. 1, pp. 111-122, 1966. [5] C. Pommerenke, "On the Hankel determinants of univalent functions," *Mathematika*, vol. 14, no. 1, pp. 108-112, 1967.
67. D. Bansal, "Upper bound of second Hankel determinant for a new class of analytic functions," *Applied Mathematics Letters*, vol. 26, no. 1, pp. 103-107, 2013.
68. M. M. Elhosh, "On the second Hankel determinant of univalent functions," *Bull. Malays. Math. Soc.*, vol. 9, no. 1, pp. 23-25, 1986.
69. J. W. Noonan and D. K. Thomas, "On the Hankel determinants of areally mean p-valent functions," *Proceedings of the London Mathematical Society*, vol. s3-25, no. 3, pp. 503-524, 1972.
70. J. W. Noonan, "Coefficient differences and Hankel determinants of areally mean p-valent functions," *Proceedings of American Mathematical Society*, vol. 46, pp. 29-37, 1974.
71. J. W. Noonan and D. K. Thomas, "On the second Hankel determinant of areally mean p-valent functions," *Transactions of the American Mathematical Society*, vol. 223, pp. 337-346, 1976.
72. M. M. Elhosh, "On the second Hankel determinant of close-toconvex functions," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 9, no. 2, pp. 67-68, 1986.
73. M. Arif, K. I. Noor, and M. Raza, "Hankel determinant problem of a subclass of analytic functions," *Journal of Inequalities and Applications*, vol. 2012, Article ID 22, 2012
74. T. Hayami and S. Owa, "Generalized Hankel determinant for certain classes," *International Journal of Mathematics and Analysis*, vol. 4, no. 49-52, pp. 2573-2585, 2010.
75. T. Hayami and S. Owa, "Applications of Hankel determinant for p-valently starlike and convex functions of order α ," *Far East Journal of Applied Mathematics*, vol. 46, no. 1, pp. 1-23, 2010.
76. T. Hayami and S. Owa, "Hankel determinant for p-valently starlike and convex functions of order α ," *General Math*, vol. 17 , no. 4 , pp. 29-44, 2009 .
77. A. Janteng, S. A. Halim, and M. Darus, "Hankel determinant for starlike and convex functions," *International Journal of Mathematics and Analysis*, vol. 1, no. 13-16, pp. 619-625, 2007.
78. A. K. Mishra and P. Gochhayat, "Second Hankel determinant for a class of analytic functions defined by fractional derivative," *International Journal of Mathematics and Mathematical Sciences*, vol. 2008, Article ID 153280, 10 pages, 2008.
79. N. Mohamed, D. Mohamad, and S. Cik Soh, "Second Hankel determinant for certain generalized classes of analytic functions," *International Journal of Mathematics and Analysis*, vol. 6, no. 17-20, pp. 807-812, 2012 .
80. G. Murugusundaramoorthy and N. Magesh, "Coefficient inequalities for certain classes of analytic functions associated with Hankel determinant," *Bulletin of Mathematical Analysis and Applications*, vol. 1, no. 3, pp. 85-89, 2009.
81. K. I. Noor, "Higher order close-to-convex functions," *Japanese Journal of Mathematics*, vol. 37, no. 1, pp. 1-8, 1992.

82. K. I. Noor, "On the Hankel determinant problem for strongly close-to-convex functions," *Journal of Natural Geometry*, vol. 11, no. 1, pp. 29-34, 1997.
83. K. I. Noor, "On certain analytic functions related with strongly close-to-convex functions," *Applied Mathematics and Computation*, vol. 197, no. 1, pp. 149-157, 2008. [23] K. I. Noor and S. A. Al-Bany, "On Bazilevic functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 10, no. 1, 88 pages, 1987.
84. K. I. Noor, "On analytic functions related with functions of bounded boundary rotation," *Commentarii Mathematici Universitatis Sancti Pauli*, vol. 30, no. 2, pp. 113-118, 1981.
85. K. I. Noor, "On meromorphic functions of bounded boundary rotation," *Caribb. J. Math.*, vol. 1, no. 3, pp. 95-103, 1982.
86. K. I. Noor, "Hankel determinant problem for the class of functions with bounded boundary rotation," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 28, no. 8, pp. 731-739, 1983.
87. K. I. Noor and I. M. A. Al-Naggar, "On the Hankel determinant problem," *Journal of Natural Geometry*, vol. 14, no. 2, pp. 133-140, 1998.
88. R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "The Fekete-Szegő coefficient functional for transforms of analytic functions," *Bulletin of the Iranian Mathematical Society*, vol. 35, no. 2, pp. 119-142, 2009.
89. R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Coefficient bounds for p_- -valent functions," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 35-46, 2007. W. C. Ma and D. Minda, "A united treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis*, Tianjin, 1992.
90. R. K. Raina and J. Sokół, "On coefficient estimates for a certain class of starlike functions, Hacet," *Journal of Mathematics and Statistics*, vol. 44, no. 6, pp. 1427-1433, 2015.
91. G. Murugusundaramoorthy and T. Bulboacă, "Hankel determinants for new subclasses of analytic functions related to a shell shaped region," *Mathematics*, vol. 8, no. 6, article 1041, 2020.
92. S. Gandhi, "Radius Estimates for Three Leaf Function and Convex Combination of Starlike Functions," in *Mathematical Analysis I: Approximation Theory. ICRAPAM 2018*, vol. 306 of *Springer Proceedings in Mathematics & Statistics*, Springer, Singapore.
93. S. N. Malik, S. Mahmood, M. Raza, S. Farman, and S. Zainab, "Coefficient inequalities of functions associated with petal type domains," *Mathematics*, vol. 6, no. 12, p. 298, 2018.
94. K. Sharma, N. K. Jain, and V. Ravichandran, "Starlike functions associated with a cardioid," *Afrika Matematika*, vol. 27, pp. 923-939, 2016.
95. V. S. Masih and S. Kanas, "Subclasses of starlike and convex functions associated with the Limaçon domain," *Symmetry*, vol. 12, no. 6, p. 942, 2020.
96. Y. Yuzaimi, A. H. Suzeini, and A. B. Akbarally, "Subclass of starlike functions associated with a limaçon," *AIP Conference Proceedings*, vol. 1974, article 030023, 2018.
97. P. L. Duren, *Univalent Functions. Grundlehren der Mathematischen Wissenschaften*, vol. 259, Springer, New York, 1983.
98. U. Grenander and G. Szegő, *Toeplitz Forms and Their Applications*. California Monographs in Mathematical Sciences, University of California Press, Berkeley, 1958.
99. M. Abramowitz and I. A. Stegun (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover Publications Inc., New York, 1965.
100. A. A. Attiya, "Some applications of Mittag-Leffler function in the unit disk," *Filomat* 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>

101. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, *Complex Var. Elliptic Equ.* 61(3) (2016), 338-350.
<https://doi.org/10.1080/17476933.2015.1079628>
102. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, *Commun. Korean Math. Soc.* 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
103. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, *Fract. Calc. Appl. Anal.* 5(3) (2002), 303-313.
104. P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften Series 259, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, *Kyungpook Math. J.* **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>
105. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, *Konuralp J. Math.* 7(1) (2019), 199-202.
106. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, *Stud. Univ. Babeş-Bolyai Math.* 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbmath.2020.1.06>
107. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, *Afr. Mat.* 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
108. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, *J. Appl. Math.* 2011(2011), Article ID 298628.
<https://doi.org/10.1155/2011/298628>
109. V. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.
110. E. Merkes and B. T. Scott, Starlike hypergeometric functions, *Proc. Amer. Math. Soc.* 12 (1961), 885-888.
111. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, *C. R. Acad. Sci. Paris* 137(1903), 554 – 558.
112. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, *Integral Transforms Spec. Funct.* 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
113. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, *Novi Sad J. Math.* **49**(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
114. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, *Hacet. J. Math. Stat.* 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
115. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, *Afr. Mat.* 28(2017), 1357-1366.
<https://doi.org/10.1007/s13370-017-0520-x>
116. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, *Afr. Mat.* 27(5) (2016), 1021-1027.
<https://doi.org/10.1007/s13370-016-0398-z>
117. M. S. Robertson, On the theory of univalent functions, *Ann. of Math.* (2) 37(2) (1936), 374-408.

118. Salah, Jamal, Hameed Ur Rehman, and Iman Al-Buwaiqi. "The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function." *Mathematics and Statistics* 10.2 (2022): 410-418.
119. Rehman, Hameed Ur, Maslina Darus, and Jamal Salah. "Graphing Examples of Starlike and Convex Functions of order β ." *Appl. Math. Inf. Sci* 12.3 (2018): 509-515.
120. Salah, Jamal Y., and O. M. A. N. Ibra. "Properties of the Modified Caputo's Derivative Operator for certain analytic functions." *International Journal of Pure and Applied Mathematics* 109.3 (2014): 665-671.
121. Jamal Salah and Maslina Darus, On convexity of the general integral operators, An. Univ. Vest Timis. Ser. Mat. -Inform. 49(1) (2011), 117-124.
122. Salah, Jamal. "TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS." (2019).
123. H. Silverman, Starlike and convexity properties for hypergeometric functions, J. Math. Anal. Appl. 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
124. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, Integral Transforms Spec. Funct. 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
125. L. Spáček, Contribution à la théorie des fonctions univalentes, Časopis Pro Pěstování Matematiky **62**(1932), 12 – 19.
126. Salah, Jamal Y. "A new subclass of univalent functions defined by the means of Jamal operator." *Far East Journal of Mathematical Sciences (FJMS)* Vol 108 (2018): 389-399.
127. Jamal Y. Salah On Riemann Hypothesis and Robin's Inequality. International Journal of Scientific and Innovative Mathematical Research (IJSIMR). Volume (3) 4 (2015) 9-14.
128. Salah, Jamal. "Neighborhood of a certain family of multivalent functions with negative coefficients." *Int. J. Pure Appl. Math* 92.4 (2014): 591-597.
129. Salah, Jamal, and Maslina Darus. "A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional." *J. Appl. comp Sc. Math* 5.1 (2011): 97-101.
130. J. Salah, Certain subclass of analytic functions associated with fractional calculus operator, Trans. J. Math. Mech., 3 (2011), 35–42. Available from: <http://tjmm.edyopress.ro/journal/11030106.pdf>.
131. Salah, Jamal. "Some Remarks and Propositions on Riemann Hypothesis." *Mathematics and Statistics* 9.2 (2021): 159-165.
132. Salah, Jamal Y. Mohammad. "The consequence of the analytic continuity of Zeta function subject to an additional term and a justification of the location of the non-trivial zeros." *International Journal of Science and Research (IJSR)* 9.3 (2020): 1566-1569.
- 133.
134. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, Journal of Inequalities in Pure and Applied Mathematics 5(4) (2004), Article ID 83, 10 pages.
135. A. Wiman, Über die nullstellen der funktionen $E_{\alpha}(x)$, Acta Math. 29 (1905), 217-134.

136. Salah, Jamal Y. Mohammad. "An Alternative perspective to Riemann Hypothesis." *PSYCHOLOGY AND EDUCATION* 57.9 (2020): 1278-1281.
137. Salah, Jamal Y. "A note on the Hurwitz zeta function." *Far East Journal of Mathematical Sciences (FJMS)* 101.12 (2017): 2677-2683.
138. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. *Far East Journal of Mathematical Sciences (FJMS)*. Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
139. Jamal Y. Salah, A Note on Riemann Zeta Function, *IOSR Journal of Engineering (IOSRJEN)*, vol. 06, no. 02, pp. 07-16, February 2016, URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
140. Jamal Y. Salah, A Note on Gamma Function, *International Journal of Modern Sciences and Engineering Technology (IJMSET)*, vol. 2, no. 8, pp. 58-64, 2015.
141. Salah, Jamal, and S. Venkatesh. "Inequalities on the Theory of Univalent Functions." *Journal of Mathematics and System Science* 4.7 (2014).
142. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, *Yokohama Math. J.* 19 (1971), 7 – 15.
143. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, *Progr. Fract. Differ. Appl.* Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL:
144. Jamal Salah. Subordination and superordination involving certain fractional operator. *Asian Journal of Fuzzy and Applied Mathematics*, vol. 1, pp. 98-107, 2013. URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>
145. Salah, Jamal Y. Mohammad. "Two Conditional proofs of Riemann Hypothesis." *International Journal of Sciences: Basic and Applied Research (IJSBAR)* 49.1 (2020): 74-83.
146. Salah, Jamal. "Fekete-Szego problems involving certain integral operator." *International Journal of Mathematics Trends and Technology-IJMTT* 7 (2014).